Board Logo

ANAPHYLAXIS
Plan of Care (Sample)
STUDENT INFORMATION

Student Name	Date Of Birth	Insert Photo
Ontario Ed. #	Age	
Grade	Teacher(s)	
	Medical ID jewellery 🛛 Yes 🗌 No	

EMERGENCY CONTACTS (LIST IN PRIORITY)						
NAME	RELATIONSHIP	DAYTIME PHONE	ALTERNATE PHONE			
1.						
2.						
3.						

KNOWN LIFE-THREATENING TRIGGERS			
CHECK (✓) THE APPROPRIATE BOXES			
Food(s): Insect Stings:			
□ Other:			
Epinephrine auto-injector(s) expiry date(s):			
Dosage: EpiPen Jr [®] EpiPen [®] 0.15 mg 0.3 mg			
Previous anaphylactic reaction: Student is at greater risk.			
☐ Has asthma. Student is at greater risk . If student is having a reaction and has difficulty breathing, give epinephrine before asthma medication.			
Any other medical condition or allergy?			
Page 1 of 4			

DAILY/ROUTINE ANAPHYLAXIS MANAGEMENT

SYMPTOMS

A STUDENT HAVING AN ANAPHYLACTIC REACTION MIGHT HAVE <u>ANY</u> OF THESE SIGNS AND SYMPTOMS:

- Skin system: hives, swelling (face, lips, tongue), itching, warmth, redness.
- **Respiratory system** (breathing): coughing, wheezing, shortness of breath, chest pain or tightness, throat tightness, hoarse voice, nasal congestion or hay fever-like symptoms (runny, itchy nose and watery eyes, sneezing), trouble swallowing.
- Gastrointestinal system (stomach): nausea, pain or cramps, vomiting, diarrhea.
- **Cardiovascular system** (heart): paler than normal skin colour/blue colour, weak pulse, passing out, dizziness or lightheadedness, shock.
- **Other**: anxiety, sense of doom (the feeling that something bad is about to happen), headache, uterine cramps, metallic taste.

EARLY RECOGNITION OF SYMPTOMS AND IMMEDIATE TREATMENT COULD SAVE A PERSON'S LIFE.

Avoidance of an allergen is the main way to prevent an allergic reaction.

Food Allergen(s): (The amount required to cause a reaction varies by person and in some people, it can be triggered by a small amount.)

Food(s) to be avoided:

Safety measures: _____

Insect Stings: (Risk of insect stings is higher in warmer months. Avoid areas where stinging insects nest or congregate. Destroy or remove nests, cover or move trash cans, keep food indoors.)

Designated eating area inside school building _____

Safety measures:

Other information:

Page 2 of 4

DICATION (Epinephrine auto-injectors):
cess to epinephrine auto-injector: ident requires assistance to access their auto-injector?
cation: With:
ner:
o, student will carry their auto-injector at all times: in the classroom, outside the classroom (e.g., ary, cafeteria/lunchroom, gym) and off-site (e.g., field trips/ excursions).
to-injector in student's: Backpack/fanny pack Other (specify)
ditional auto-injector: e student has an additional auto-injector at school? □ Yes □ No es, the additional auto-injector is kept:
cation: With: With:
ner:
EMERGENCY PROCEDURES
(DEALING WITH AN ANAPHYLACTIC REACTION)
(DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN
(DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN
(DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN ET WORSE QUICKLY. TEPS 1. Give epinephrine auto-injector (e.g. EpiPen [®]) at the first sign of a known or suspected
(DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN ET WORSE QUICKLY. TEPS 1. Give epinephrine auto-injector (e.g. EpiPen [®]) at the first sign of a known or suspected anaphylactic reaction.
 (DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN ET WORSE QUICKLY. TEPS Give epinephrine auto-injector (e.g. EpiPen[®]) at the first sign of a known or suspected anaphylactic reaction. Call 9-1-1. Tell them someone is having a life-threatening allergic reaction. Give a second dose of epinephrine as early as five (5) minutes after the first dose if there
 (DEALING WITH AN ANAPHYLACTIC REACTION) CT QUICKLY. THE FIRST SIGNS OF A REACTION CAN BE MILD, BUT SYMPTOMS CAN ET WORSE QUICKLY. TEPS Give epinephrine auto-injector (e.g. EpiPen[®]) at the first sign of a known or suspected anaphylactic reaction. Call 9-1-1. Tell them someone is having a life-threatening allergic reaction. Give a second dose of epinephrine as early as five (5) minutes after the first dose if there is no improvement in symptoms. Follow direction of emergency personnel, including transport to hospital (ideally by ambulance), even if symptoms are mild or have stopped. The reaction could worsen or come back, even after treatment. Stay in the hospital for an appropriate period of observation as decided by the emergency department physician (generally about 4 - 6

HEALTHO	HEALTHCARE PROVIDER INFORMATION (OPTIONAL)				
Healthcare provider may in Respiratory Therapist, Cert			oner, Registered Nurse, Pharmacist, ified Asthma Educator.		
Healthcare Provider's Name	e:				
Profession/Role:					
Signature:		Date:			
Special Instructions/Notes/F					
for which the authorization t	o administer applie	es, and possible			
★This information may rem	ain on file if there a	ire no changes	to the student's medical condition.		
	AUTHORIZAT	ION/PLAN R	EVIEW		
INDIVIDUALS	WITH WHOM TH	IS PLAN OF CA	ARE IS TO BE SHARED		
1	2		3		
4			6		
Other individuals to be cont Before-School Program	acted regarding Pla □Yes	an Of Care: No			
After-School Program	☐ Yes	🗌 No			
School Bus Driver/Route #	(If Applicable)				
Other:					
This plan remains in effec	t for the 20	20 school y	year without change and will be (It is the parent(s)/guardian(s) e the plan of care during the school		
Parent(s)/Guardian(s):	Signatura		_ Date:		
Student:	Signature		_ Date:		
Principal:					